

University of Dundee

An Aphid Effector Targets Trafficking Protein VPS52 in a Host-Specific Manner to Promote Virulence

Rodriguez, Patricia A.; Escudero Martinez, Carmen; Bos, Jorunn

Published in:
Plant Physiology

DOI:
[10.1104/pp.16.01458](https://doi.org/10.1104/pp.16.01458)

Publication date:
2017

Document Version
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Rodriguez, P. A., Escudero Martinez, C., & Bos, J. (2017). An Aphid Effector Targets Trafficking Protein VPS52 in a Host-Specific Manner to Promote Virulence. *Plant Physiology*, 173(3), 1892-1903.
<https://doi.org/10.1104/pp.16.01458>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

An aphid effector targets trafficking protein VPS52 in a host-specific manner to promote virulence

Rodriguez, Patricia A.², Escudero-Martinez, Carmen^{1,2}, and Bos, Jorunn I. B.^{1,2}

¹Cell and Molecular Sciences, The James Hutton Institute, Dundee, DD2 5DA, UK.

²Division of Plant Sciences, School of Life Sciences, University of Dundee, Dundee DD2 5DA, UK.

One sentence summary: A secreted salivary protein from an herbivorous insect targets a protein in host plants to promote infestation

1

¹ List of author contributions

P.R. and J.B. designed the experiments. P.R., C.E-M, and J.B. performed the experiments and analyzed the data. P.R. and J.B. wrote the manuscript. J.B. directed the project. All authors read and approved the final manuscript.

Funding information: Research was supported the Biotechnology and Biological Science Research Council (BB/J005258/1, to J.I.B.B.), the European Research Council (APHIDHOST-310190 to J.I.B.B), and the Royal Society of Edinburgh (Personal Research Fellowship, to J.I.B.B.).

Corresponding author:

Jorunn I. B. Bos,
Division of Plant Sciences, School of Life Sciences, University of Dundee
Cell and Molecular Sciences, The James Hutton Institute
Dundee, DD2 5DA, UK.
j.bos@dundee.ac.uk

Abstract

Plant- and animal-feeding insects secrete saliva inside their hosts, containing effectors, which may promote nutrient release and suppress immunity. Although for plant pathogenic microbes it is well established that effectors target host proteins to modulate host cell processes and promote disease, the host cell targets of herbivorous insects remain elusive. Here, we show that the existing plant pathogenic microbe effector paradigm can be extended to herbivorous insects in that effector-target interactions inside host cells modify critical host processes to promote plant susceptibility. We showed that the effector Mp1 from *Myzus persicae* associates with the host Vacuolar-Protein-Sorting-Associated Protein 52 (VPS52). Using natural variants we provide a strong link between effector virulence activity and association with VPS52, and show that the association is highly specific to *Myzus persicae*-host interactions. Also, co-expression of Mp1, but not Mp1-like variants, specifically with host VPS52s resulted in effector relocalization to vesicle-like structures that associate with prevacuolar compartments. We show that high VPS52 levels negatively impact virulence, and that aphids are able to reduce VPS52 levels during infestation, indicating that VPS52 is an important virulence target. Our work is an important step forward in understanding, at the molecular level, how a major agricultural pest promotes susceptibility during infestation of crop plants. We give evidence that an herbivorous insect employs effectors that interact with host proteins as part of an effective virulence strategy, and that these effectors likely function in a species-specific manner.

Introduction

Many insect species secrete saliva, containing various proteins, inside their host to enable feeding. The identification and characterization of insect salivary molecules over recent years has contributed to novel insights into suppression of host immune responses. Aphids need to form a close association with their host to enable feeding and infestation. These insects use specialized mouthparts, or stylets, to penetrate the leaf surface and establish a feeding site. While probing and feeding, aphids secrete saliva into different host cell types as well as the apoplast¹. Advances in genomics and proteomics have facilitated the identification of a diverse array of proteins in aphid saliva as well as aphid salivary glands (Harmel et al., 2008; Carolan et al., 2009; Bos et al., 2010). These include proteins with predicted functions, such as metalloproteases, disulfide isomerases, calreticulins, ARMET proteins, glutathione peroxidases, and CLP-domain serine proteases.

54 However, nearly half of the predicted aphid salivary proteins described have no functional
55 annotation and/or similarity to proteins in other organisms (Rodriguez and Bos, 2013). The
56 abundance though of these proteins in aphid saliva/salivary glands, suggests they exhibit
57 an important effector activity towards promoting aphid virulence.

58
59 Upon interaction with plant pathogen microbes, most plants are able to defend themselves
60 by recognizing conserved parasite molecules, or PAMPs (Pathogen-Associated-Molecular-
61 Pattern), to activate PTI (PAMP-Triggered-Immunity). However, successful plant
62 pathogenic microbes deliver effectors inside their hosts to suppress this and other types of
63 plant defenses by interacting with and altering the mode-of action of important plant
64 defense signaling components. Over the past decade studies on plant microbe effectors
65 and their activities has revealed exciting insight into the host cell processes targeted to
66 enable infection. For example, effectors from oomycete plant pathogens have been shown
67 to target an ubiquitin E3 ligase (Bos et al., 2010), a MAPKKK (King et al., 2014), a host
68 autophagosome protein (Dagdas et al., 2016) and a host phosphatase (Boevink et al.,
69 2016) to suppress host immunity. Other host targets include JAZ proteins involved in JA-
70 signalling, which are targeted by both bacterial and fungal effectors (Jiang et al., 2013;
71 Plett et al., 2014), but also the host proteasome (Groll et al., 2008; Ustun et al., 2013),
72 extracellular proteases (Song et al., 2009) as well as the cytoskeleton and secretion
73 pathways (Bozkurt et al., 2011; Lee et al., 2012). Also aphids alter host physiology as
74 reflected by their ability to affect nutrient allocation (Sandström et al., 2000; Gironse et al.,
75 2005) and to suppress defence responses (Bos et al., 2010; Elzinga and Jander, 2013;
76 Rodriguez et al., 2014). With parallels emerging between plant microbe infection and
77 aphid infestation strategies, it is likely that aphid effectors also target important host cell
78 processes to promote virulence or effector-triggered-susceptibility (ETS) (Rodriguez and
79 Bos, 2013).

80
81 Although there is limited insight into the molecular basis of plant pathogen and pest host
82 range, effectors and their plant targets are predicted to be involved. For example, an
83 effector may only be able to interact with a specific plant protein in host but not nonhost
84 plants to suppress defenses as a consequence of target diversification (Schulze-Lefert
85 and Panstruga, 2011). Indeed, Zheng and co-authors (Zheng et al., 2014) identified 8
86 *Phytophthora infestans* effectors that were able to suppress flg22-activated reporters, of
87 which 5 were only able to do so in host tomato but not in nonhost *Arabidopsis* protoplasts.
88 Among the different aphid species, *Myzus persicae*, is one of the major pests, which is

partly due to its broad host range, which includes plants in over 40 families. How this species is able to infest such a wide range of plant species and whether this involves secretion of effectors remains to be elucidated.

To date a number of aphid effectors has been identified that affect aphid virulence as determined by *in planta* over-expression assays as well as RNAi in aphids (Mutti et al., 2008; Bos et al., 2010; Atamian et al., 2013; Pitino and Hogenhout, 2013; Elzinga et al., 2014). This includes effectors from the broad host range aphid species *M. persicae* that reduce aphid virulence upon over-expression, including Mp10, Mp42, Mp56, Mp57, and Mp58 (Bos et al., 2010; Elzinga et al., 2014). Importantly, several aphid effectors have been identified to date that contribute to aphid virulence. Effector C002, first identified in the *Acyrtosiphum pisum* (pea aphid), contributes to aphid survival as evidenced by RNAi experiments that resulted in a reduction of C002 transcript levels and reduced aphid virulence (Mutti et al., 2008; Pitino and Hogenhout, 2013). Moreover, overexpression of *M. persicae* C002 in *Arabidopsis* and *Nicotiana benthamiana* enhances *M. persicae* virulence. Other effectors found to enhance aphid virulence upon over-expression in host plants are Me10 and Me23 from *Macrosiphum euphorbiae* (potato aphid), as well as Mp1 (PIntO1), Mp2 (PIntO2) and Mp55 from *M. persicae* (Atamian et al., 2013; Pitino and Hogenhout, 2013; Elzinga et al., 2014). For several of these effectors, including Mp1, there is strong evidence for secretion into the plant-host interface, as proteomics-based approaches revealed their presence in aphid saliva (Harmel et al., 2008; Carolan et al., 2009; Carolan et al., 2011). Another interesting observation is that there is specificity with regards to activity of similar effectors from different species (Pitino and Hogenhout, 2013; Elzinga and Jander, 2014). While Mp1 promotes *M. persicae* virulence upon over-expression in the phloem of *Arabidopsis* transgenic lines, Ap1, an Mp1-like sequence from *A. pisum*, does not. With increasing numbers of aphid effectors being identified, the next step is to investigate their function and understand the cellular processes they target.

We performed yeast-two-hybrid (Y2H) screens against a potato library to identify aphid effector host targets and gain insight into host cellular reprogramming by aphids. We found that *M. persicae* effector Mp1 associated with Vacuolar-Protein-Sorting-Associated Protein 52 (VPS52), a component of the Golgi-Associated-Retrograde Protein (GARP) complex specifically from host but not poor-host plant species. Moreover, we implicate both the effector and host protein in host susceptibility to *M. persicae*. Our data support a model wherein aphids target a host cell trafficking pathway protein to promote infestation.

Results

Aphid effector Mp1 associates with Arabidopsis and potato VPS52

With several aphid effectors identified to date that promote virulence, including Mp1, we aimed to gain insight into the molecular mechanisms underlying virulence activity. We performed an Y2H screen against a *Solanum tuberosum* library to a depth of 7.6×10^6 yeast transformants. Yeast prey constructs were isolated from colonies recovered from selection plates, sequenced, and subjected to co-transformation with the Mp1-bait construct to identify sequences and verify interactions with individual prey plasmids. Among the prey-constructs we identified sequences corresponding to VPS52 (Vacuolar Protein Sorting-Associated Protein 52) (XP_006338692.1, 3 independent clones), FLX-like 2 (XP_015160018.1, 3 independent clones), phytochrome B (XP_006355734.1, 1 clone) and an uncharacterized protein (XP_006340727.1, 1 clone). Independent co-transformation experiments showed that yeast co-expressing a bait-Mp1 construct with a prey-StVPS52 (potato VPS52) construct was able to grow on –His medium and exhibited β -galactosidase activity (Fig. 1A). Since VPS52 has been previously identified and characterized in *Arabidopsis* (Lobstein et al., 2004; Guermonprez et al., 2008), also a host species of *M. persicae*, we tested whether Mp1 was able to interact with *Arabidopsis* VPS52 (AtVPS52) as well. AtVPS52 and StVPS52 share about 80% identity and the VPS52 sequence overall is highly conserved among plant species (Supplemental Fig. 1). Yeast reporter assays showed that Mp1 was able to interact with both AtVPS52 and StVPS52 (Fig. 1A).

To independently confirm the association of VPS52 with Mp1 *in planta*, we cloned StVPS52 and AtVPS52 into expression vectors with N-terminal epitope tags, as well as Ap1, an Mp1-like effector from the aphid species *A. pisum* (pea aphid). Co-expression of GFP-StVPS52 or GFP-AtVPS52 with FLAG-Mp1 or FLAG-Ap1 in *N. benthamiana*, followed by immunoprecipitation using GFP-trap beads confirmed that only Mp1 associated with both AtVPS52 and StVPS52 inside plant cells (Fig. 1B and C). This correlates with the ability of Mp1, but not Ap1, to promote *M. persicae* virulence on *Arabidopsis* (Pitino and Hogenhout, 2013).

We cloned a set of additional Mp1-like sequences from the aphid species *Rhopalosiphum padi* (bird cherry-oat aphid), and *Myzus cerasi* (black cherry aphid) to determine whether Mp1-like effectors from additional species limited in their ability/unable to infest potato or *Arabidopsis* were also able to associate with AtVPS52 or StVPS52 (Supplemental Fig. 2). Co-immunoprecipitations with these Mp1-like effectors were performed in parallel with Mp1 to test for association with VPS52. Although we occasionally detected faint bands for Mp1-like variants on western blots in the GFP or GFP-VPS52 pull-downs, which likely reflects nonspecific binding to the GFP-magnetic beads, we only consistently detected a strong band corresponding to Mp1 in the GFP-VPS52 pull downs (Fig. 2A). Based on this, we conclude that none of the Mp1-like effectors were pulled down by GFP-VPS52 despite showing comparable expression levels to Mp1 (Fig. 2A). This provides further evidence that the association of Mp1 with VPS52 is specific to Mp1, but not Mp1-like effectors.

The Mp1-VPS52 association takes place in a host-specific manner

Although *M. persicae* has a broad host range, this aphid shows poor performance on *Medicago truncatula* and barley (*Hordeum vulgare*) under controlled conditions, and does not cause significant infestations on these plants in a natural environment (Gao et al., 2007; Davis and Radcliffe, 2008). To determine whether Mp1 is able to associate with VPS52 proteins from poor-host plant species of *M. persicae*, we cloned VPS52 sequences from barley and *M. truncatula* into plant expression vectors. We performed co-expression experiments of GFP-VPS52s and FLAG-Mp1 in *Nicotiana benthamiana* followed by co-immunoprecipitation. Of the different VPS52 proteins, StVPS52 showed the lowest expression level in the input as determined by western blotting. Despite this, only StVPS52 and AtVPS52, but not HvVPS52 and MtVPS52, were able to pull-down Mp1 (Fig. 2B). Also, we noted that Mp1 protein levels in the input were more abundant in the presence of StVPS52 and AtVPS52, but not HvVPS52 and MtVPS52, indicating that the association may stabilize the effector protein (Fig. 2B).

Since *R. padi* and *A. pisum* are able to infest barley and *M. truncatula*, respectively, we tested whether the Mp1-like proteins from these aphid species were able to associate with HvVPS52 and MtVPS52. Co-expression in *N. benthamiana* followed by co-immunoprecipitation showed that GFP-HvVPS52 and -MtVPS52 did not pull down any of the Mp1-like proteins in these experiments (Supplemental Fig. 3SA and B). This suggests that the Mp1-VPS52 association may be specific to *M. persicae* and its host plants.

Co-expression of Mp1 with AtVPS52 or StVPS52 results in its relocalization to vesicle-like structures that associate with prevacuolar compartments

In plants, VPS52 has been previously identified and characterized in *Arabidopsis* (Lobstein et al., 2004; Guermonprez et al., 2008), where this protein was shown to localize partially to post-Golgi and prevacuolar compartments suggesting it may have similar functions in trafficking as in other eukaryotes. We further investigated the (co-)localisation of VPS52 and Mp1 upon overexpression in *N. benthamiana* and found that StVPS52, similar to GFP-AtVPS52, localized in mobile vesicle-like structures, which partially co-localized with the prevacuolar compartment marker PS1 (Kotzer et al., 2004) (Fig. 3A and B). Limited co-localization was observed with the endomembrane trafficking compartment markers Ara6 and Ara7 (Supplemental Fig. 4). In the absence of over-expressed VPS52, GFP-Mp1 (± 35 kDa) predominantly localized to the cytoplasm and nucleus similar to the free GFP control (Fig. 3C). However, in the presence of over-expressed mRFP-AtVPS52 or -StVPS52, Mp1 re-localised to vesicle-like structures that were associated with RFP-VPS52 (Figure 3C). Similar to StVPS52 and AtVPS52, MtVPS52 and HvVPS52 localized to vesicle-like structures upon over-expression in *N. benthamiana* (Supplemental Fig. 5 and 6). However, no Mp1-relocalization was observed upon co-expression with HvVPS52 or MtVPS52 (Supplemental Fig. 5), indicating that relocalization only takes place when Mp1 is co-expressed with VPS52s that are able to associate with it.

In addition, we performed co-localization of the Mp1-like effectors from *A. pisum* (Ap1), *R. padi* (Rp1) and *M. cerasi* (Mc1) with the VPS52s cloned from the 4 different plant species. GFP-Ap1, -Mc1 and -Rp1 were only detected in the cytoplasm and nucleus, similar to free GFP, in the absence of over-expressed VPS52 (Supplemental Fig. 6). Western blotting confirmed the different GFP-Mp1-like fusions were expressed as full-length proteins (Supplemental Fig. 7). No relocalization was observed with any of the Mp1-like effector-VPS52 combinations. These data are in line with the co-immunoprecipitation data and provide further evidence that the Mp1-VPS52 association may be species specific.

Phloem-specific over-expression of Mp1, but not Mp1-like effectors from other species, promotes *Myzus persicae* virulence in *Nicotiana benthamiana*

Whilst Mp1, but not Ap1, promotes virulence when expressed in the phloem companion cells of *Arabidopsis* transgenic lines (Pitino and Hogenhout, 2013), no virulence activity

was found in *N. benthamiana* using 35S-based expression in leaf discs (Bos et al., 2010). We used transient assays in intact *N. benthamiana* plants, similar to Elzinga et al. (2014), to determine whether Mp1 and Mp1-like effectors impact aphid virulence when expressed in the phloem companion cells of a solanaceous host. We expressed Mp1, Ap1, Rp1 and Mc1 under the control of the AtSUC2 phloem-specific promoter and challenged infiltrated leaf areas with *M. persicae* 1st-instar nymphs. Virulence, as measured by nymph production, was monitored over a 14-day period, with aphids being moved to fresh infiltration sites every 6 days. While expression of Mp1 resulted in a 64% increase of *M. persicae* nymph production compared to the vector control (Fig. 4), none of the Mp1-like effectors showed any significant effect on virulence. The different Mp1-like effectors were all equally stable in *planta* when expressed as GFP-, FLAG- and Myc-fusion proteins, suggesting that the observed differences in virulence activity are not due to differences in protein stability (Fig. 1, Supplemental Fig. 6, Supplemental Fig. 7). Importantly, the lack of virulence activity of the Mp1-like effectors from different aphid species towards *M. persicae* correlates with the inability to associate with StVPS52 or AtVPS52.

Phloem-specific over-expression StVPS52 reduces *Myzus persicae* virulence

We were interested to determine the impact of VPS52 on aphid virulence and tested whether we could use Virus-Induced-Gene-Silencing (VIGS) of VPS52 in *N. benthamiana* in combination with *M. persicae* performance assays. Aphids were unable to survive on control plants in any attempted VIGS experiment, suggesting that, in our hands, VIGS was incompatible with *M. persicae* performance assays. We then decided to over-express StVPS52 in a solanaceous host, *N. benthamiana*, under a phloem-specific promoter to determine whether this affected aphid virulence. Over-expression of StVPS52 reduced nymph production by about 40% compared to the vector control (Fig. 5). This suggests that high levels of host StVPS52 in phloem companion cells negatively impact *M. persicae* virulence.

AtVPS52 and StVPS52 protein levels are reduced upon aphid infestation

To investigate whether VPS52 is targeted by aphids during infestation, we made use of an available *Arabidopsis* mutant line, *pok* (Lobstein et al., 2004), which harbours a T-DNA GUS insertion in the 10th exon resulting in an active translational fusion. Homozygous plants cannot be obtained for this mutant due to a defect in male gametophyte formation

(Lobstein et al., 2004). qRT-PCR analyses of *AtVPS52* transcripts in the hemizygous mutant, did not show any differences in transcript levels indicating that there is no reduction of expression in this line (Supplemental Fig. 8A). Also, aphid fecundity experiments did not show any significant difference in aphid susceptibility of this line compared to Ws-4 wild-type plants (control) (Supplemental Fig. 8B). Using GUS-staining we assessed expression of the translational VPS52-GUS fusion in both uninfested and aphid-infested plants. Although no VPS52-GUS expression was detected in rosette leaves, we found high levels of *AtVPS52*-GUS in apical stem tissues of flowering plants (Fig. 6A, left panels) and, as previously described, in flower tissues (Guermonprez et al., 2008). The high level of VPS52 in these tissues may reflect increased vesicle trafficking taking place to allow for growth and development. Interestingly, aphid infestation strongly reduced *AtVPS52*-GUS in the stem and flower tissues (Fig. 6A, right panels). However, no reduction of *AtVPS52* transcript levels was detected by qRT-PCR upon aphid infestation suggesting the aphids target VPS52 at the post-translational level (Supplemental Fig. 8C).

Using transient-overexpression of GFP-StVPS52 or GFP-*AtVPS52* in *N. benthamiana* followed by *M. persicae* challenge, we independently confirmed the aphid-mediated reduction of VPS52, but not free GFP, at the post-translational level (Fig. 6B). In parallel, we tested whether PAMP treatment or infection with a plant pathogen similarly affected GFP-VPS52 stability in *N. benthamiana*. We found that neither the bacterial PAMP flg22 nor the oomycete plant pathogen *Phytophthora capsici* reduced GFP-StVPS52 protein levels (Fig. 6C). In contrast, a slight increase of StVPS52 was detected when infiltrated tissues were exposed to flg22 or *P. capsici*. Therefore, the strong reduction of VPS52 levels specifically by *M. persicae* is unlikely the result of activation or suppression of a general plant defence response. Co-expression assays of VPS52 with Mp1 (Fig. 6B, and Fig. 1B, C, and D) did not show any evidence of a role for Mp1 in VPS52 degradation, suggesting that additional effectors in aphid saliva may be involved in the targeting and degradation of VPS52. Since phloem-specific over-expression of VPS52 negatively impacted aphid virulence, aphid-mediated degradation of this protein may be an important step during infestation.

During aphid infestation experiments on flowering *Arabidopsis* plants, we noticed that aphids were more abundant on stems and flower tissues than on rosette leaves of flowering plants. We therefore, followed this up with aphid choice-experiments, where 30 alate *M. persicae* aphids were released in a cage containing 4 flowering plants. We

monitored aphid numbers after 12 days on rosette leaves or stems plus flowers. This showed that aphid colonization predominantly occurred on the stems and flowers as opposed to the rosette leaves, with around 6 times more aphids being present in these tissues (Fig. 7). The high level of VPS52 expression in these aphid-preferred tissues suggests that aphids indeed need to target this protein as part of an effective infestation strategy.

Discussion

Here, we report the targeting of a host cell trafficking protein, VPS52, by an herbivorous insect. We used an aphid effector, Mp1, which promotes virulence, as a probe to identify potential aphid targets in host plants. We showed that Mp1 specifically associates with VPS52 from several host but not poor-host plant species. Moreover, we were unable to detect associations of Mp1-like effectors from other aphid species with host VPS52s suggesting that the Mp1-VPS52 association is specific to *M. persicae* and its hosts. Over-expression of StVPS52 in a solanaceous host reduced *M. persicae* performance indicating VPS52 is likely an important virulence target. Overall, our work provides important evidence that herbivorous insects, similar to plant pathogenic microbes, secrete effectors inside host cells that interact with host proteins and modify their activity to facilitate infestation.

The association of Mp1 with VPS52 was highly specific in that only Mp1, but not Mp1-like effectors from other aphid species, interacted with VPS52 from *M. persicae* hosts. Since the host species *Arabidopsis* and potato are in different families, *M. persicae* effector Mp1 may have evolved to associate with the same host protein in at least two distantly related plant species. The VPS52 family in plants is highly conserved, suggesting that perhaps minor structural differences among its members have a significant impact on the association with Mp1. By including both different variants of Mp1 and VPS52 from different aphid and plant species, we were able to show that Mp1 association with VPS52 is linked to virulence activity indicating this effector functions in a host species-specific manner. Host-specific effector functions have been proposed by Pitino and Hogenhout (2013) (Pitino and Hogenhout, 2013), who showed that *M. persicae* effectors C002, PlntO1 (Mp1), and PlntO2 (Mp2), but not their putative orthologs from the pea aphid, promoted *M. persicae* virulence in *Arabidopsis*. Our results are in line with this and provide evidence that difference in such virulence activities are not due to differences in effector protein

333 stability. The lack of interaction between any of the Mp1-like effectors and VPS52s from
334 corresponding host plant species, suggests these effectors may have evolved to exhibit
335 different activities.

336
337 Mp1 and VPS52 virulence effects were observed upon companion-cell specific over-
338 expression of these proteins, suggesting they possibly are involved in phloem-specific
339 processes during plant-aphid interactions. When GFP is expressed under the AtSUC2
340 promoter it can pass into sieve elements and reach sink tissues, including flowers (Imlau et
341 al., 1999). Although Mp1 and VPS52 were specifically expressed in companion cells this
342 site is not necessarily the site of activity as these proteins may be localized to for example
343 the sieve elements similar to GFP. VPS52, although detected in leaves, was
344 predominantly expressed in the inflorescence stems and specific flower organs. Moreover,
345 VPS52, together with other vesicle trafficking proteins, has been detected in pumpkin
346 phloem sap by proteomics, indicating that this protein functions in the phloem (Lin et al.,
347 2009). Recent evidence suggests that membrane systems, including the Golgi apparatus
348 and the ER, may be present inside sieve elements (Lin et al., 2009; Frohlich et al., 2012).
349 Also, VPS51, another component of the GARP complex, is expressed in the plant
350 vasculature as well as flowers (Pahari et al., 2014). It is therefore possible that aphids may
351 need to manipulate host processes in specific tissues, such as the phloem.

352
353 High expression of VPS52 was detected in tissues where cell elongation takes place (i.e.
354 flower organs and apical stem), which may reflect increased vesicle trafficking to allow for
355 growth and development. Interestingly, our results show that in the case of the
356 *Arabidopsis-M. persicae* interaction, aphids prefer to feed on tissues with high levels of
357 VPS52 expression (i.e. inflorescence stems and flowers) and thus on tissues with high
358 growth or development rates. Preferred infestation by aphids, including *M. persicae*, of
359 stems, flower buds and flowers of certain host species has been reported previously
360 (Guldmond et al., 1998; Ashouri et al., 2001), and is possibly triggered by nutrient re-
361 allocation to flowers and buds during plant developmental stages. Our finding that VPS52
362 is highly expressed in these tissues and that VPS52 phloem-specific overexpression
363 negatively impacts aphid virulence suggests that aphid-mediated degradation of VPS52 is
364 important for successful infestation.

365
366 VPS52 is a component of the GARP complex, which is involved in the transport from
367 endosomes to the trans-Golgi network and has been mainly characterized in yeast and

mammalian systems (Conibear and Stevens, 2000; Conibear et al., 2003; Reggiori et al., 2003). Schindler et al. (Schindler et al., 2015) recently showed that several components of this complex, including VPS52, also take part in a complex involved in endosome to plasma membrane trafficking, pointing to different functions of GARP-complex proteins in cellular trafficking. Although the role of VPS52 in plant cellular trafficking is not well understood, this protein was previously shown to localize to post-Golgi and prevacuolar compartments suggesting it may have similar functions in trafficking as in other systems (Lobstein et al., 2004; Guermonprez et al., 2008). VPS52 interacts with at least one other component of the GARP-complex in plants, VPS51, which is involved in the maintenance of vacuolar morphology as well as leaf shape and vein patterning (Pahari et al., 2014). A recent proteomics study to identify proteins associated with endosomal and secretory pathways in *Arabidopsis* revealed the enrichment of VPS52 in pull-downs of RABF1/ARA6, RABG3f, CLC2, RABD2a/ARA5, RABF2b/ARA7 (Heard et al., 2015). This suggests that VPS52 may associate with a variety of endomembrane vesicles, perhaps through binding GTPases (Heard et al., 2015). How exactly aphids impact vesicle trafficking via VPS52 remains to be investigated and will require a better understanding of VPS52 function inside plant cells.

Our work provides important evidence that the existing plant pathogenic microbe effector paradigm can be extended to herbivorous insects in that effector-target interactions inside host cells modify critical host processes to promote plant susceptibility. The negative impact of VPS52 on aphid virulence suggests this host protein either contributes to plant immunity to aphids, or perhaps negatively regulates nutrient availability in the phloem. In recent years it has become evident that host vesicle trafficking regulates plant defence responses to pathogenic microbes (Teh and Hofius, 2014; Ben Khaled et al., 2015). For example, receptor activation, and activation of defence signalling, as well as redirecting cargo to infection sites all require host trafficking machinery. Also, plant pathogenic microbes target components of this machinery to promote disease (Nomura et al., 2006; Bozkurt et al., 2011; Gu and Innes, 2012). We showed that VPS52 degradation was specific to aphid infestation and was not observed upon activation of plant immunity and plant microbe infection. Therefore, an important next step will be to dissect how vesicle trafficking pathways mediated by VPS52 are impacted by aphid infestation and how this promotes aphid virulence. And as evidenced from our work, this would need to take into consideration activities in specific plant tissues. Ultimately, the identification and characterization of aphid effectors and their host targets will provide us with novel insights

into the virulence strategies employed by agricultural pests. With limited sources of genetic resistances available in crops a detailed understanding of these virulence strategies promises to open up new avenues for crop improvement.

Materials and Methods

Plants and growth conditions

Wild type *A. thaliana* landrace Columbia-0 (Col-0), *pok* mutants (after selection) and Ws-4 (Wassilewskija-4) were grown in Levington's M2 compost with 4 mm grit (8:1) in growth chambers after seed underwent a stratification period of 48hr at 4°C. Plants were grown under 12 hours of light per day, with a photosynthetic photon flux density of 200 $\mu\text{mol m}^{-2} \text{s}^{-1}$, at 18°C and 50% relative humidity.

Plasmid construction

The sequences of the *M. cerasi* and *R. padi* effector variants were identified by transcriptome sequencing of the different aphid species (Thorpe et al., 2016). The Mp1 (GenBank accession KY273521), Ap1 (Genbank accession KY273522), Mc1 (Genbank accession KY273524), and Rp1 (Genbank accession KY273523) coding sequences without signal peptide encoding sequences were amplified from *M. persicae*, *A. pisum*, *M. cerasi* and *R. padi* cDNA, respectively. The StVPS52 and AtVPS52 coding sequences were amplified from *S. tuberosum* and *A. thaliana*, respectively. Amplicons were cloned into the pDONR207 vector (Invitrogen, Carlsbad, CA, U.S.A) using Gateway technology. BP recombination reactions were transformed in *Escherichia coli* JM109 (Promega). Constructs were verified by sequencing. MtVPS52 (NCBI Reference Sequence: XM_003590877.2) and HvVPS52 (GenBank: AK361277.1) were synthesized and cloned into pUC57 vector and provided including suitable Gateway recombination sites by GenScript (New Jersey, USA). Subsequently, LR recombination reactions were performed using pB7WGF2 (N-terminal GFP tag), pK7WGR2 (N-terminal mRFP tag) (Karimi et al., 2002), pGWB12 (N-terminal FLAG tag) and pGWB21 (N-terminal 10xMyc tag) (Nakagawa et al., 2007) as destination vectors for transient over expression in *N. benthamiana*.

***Agrobacterium tumefaciens* infiltration assays**

Constructs were introduced into *A. tumefaciens* strain GV3101 or AGL1 by electroporation. Transformants were selected using gentamycin (12.5 $\mu\text{g/ml}$), rifampicin (50 $\mu\text{g/ml}$), and spectinomycin (50 $\mu\text{g/ml}$) for transformation into pB7WGF2, pB7WGR2 and gentamycin

(12.5 µg/ml), rifampicin (50 µg/ml), and kanamycin (50 µg/ml) for transformation into pGWB12 and pGWB21. For infiltration into leaves, recombinant strains were grown in Luria-Bertani (LB) medium with above-mentioned antibiotics, harvested, and resuspended in infiltration buffer (acetosyringone 125 µM and MgCl₂ 10mM) to reach an optical density at 600nm (OD₆₀₀) = 0.3 for western blot experiments and OD₆₀₀ = 0.1 for aphid virulence assays and confocal imaging.

Total protein extractions and co-immunoprecipitation assays

Plant tissue for co-immunoprecipitation (CO-IP) experiments was extracted with GTEN lysis buffer (10% Glycerol, 25mM Tris pH 7.5, 1mM EDTA, 150mM NaCl, 0.1% NP-40, with freshly added 10mM DTT and 1x protease inhibitor cocktail (Sigma, # P9599)). Samples were incubated for 15 min in lysis buffer at 4°C. Lysate was centrifuged at 10,000g for 10 min and the supernatant was subjected to CO-IP with GFP-Trap[®]_M agarose beads (Chromotek, # gtm-20) for affinity binding of GFP-fused proteins. Western blotting was performed with the antibodies anti-GFP (# sc-8334, Santa Cruz Biotechnology Inc, USA), anti-FLAG (# sc-166384, Santa Cruz Biotechnology Inc, USA), and anti-c-Myc (# sc-40, Santa Cruz Biotechnology Inc, USA) followed by anti-rabbit-HRP (# sc-2004, Santa Cruz Biotechnology Inc, USA), or anti-mouse-HRP (# sc-2005, Santa Cruz Biotechnology Inc, USA) to detect the corresponding epitopes. Each CO-IP experiment was repeated at least two times. Plant tissues for detecting levels of expression in total extracts were extracted using Laemmli sample buffer (4% (w/v) SDS, 20% Glycerol, 120mM Tris-Cl (pH 6.8) and 0.02% (w/v) bromophenol blue) and then subjected to western blotting procedures.

flg22 and *P. capsici* infection assays

Nicotiana benthamiana leaves expressing GFP-StVPS52 2 days after agroinfiltration were drop-inoculated with two 10µl droplets of *P. capsici* LT1534 zoospores in water (500,000 spores/ml) or water (control). Tissue was collected 24hr later and was used for western blot experiments. For flg22 elicitor assay, *N. benthamiana* leaves expressing GFP-StVPS52 2 days after agroinfiltration were syringe infiltrated with, 1mM flg22 and tissue was harvested after 3hr of elicitor treatment.

***Myzus persicae* rearing**

Aphid experiments were done with *M. persicae* individuals, genotype O, reared on oil seed rape (*Brassica napus*) plants, with a long day (12h) day light and a photosynthetic photon flux density of 200 $\mu\text{mol m}^{-2} \text{s}^{-1}$ at 18°C and 50% relative humidity.

Aphid virulence assays on whole plants

For whole plant assays, *N. benthamiana* leaves were infiltrated with recombinant *A. tumefaciens* GV3101 carrying constructs to express the effectors (Mp1, Ap1, Mc1 and Rp1) or StVPS52 under the AtSUC2-promoter (Gottwald et al., 2000) at an OD₆₀₀ of 0.1. Adult aphids were placed on the underside of leaves one day after agroinfiltration using clip cages. The next day, adult aphids were removed and 3 1-st instar nymphs were left per infiltration site inside of a clip cage. Seven days after agroinfiltration, nymphs were transferred to newly infiltrated leaves, and aphid progeny was counted after 14 days of initial agroinfiltration. For aphid reproduction assays on *Arabidopsis pok* mutants and Ws-4 wild-type plants, mutant plants were firstly selected on Murashige-Skoog (MS) medium containing kanamycin. Seedlings were moved and potted into arabidopsis mix, and after 2 weeks under long-day conditions, aphid assays were performed. Two adult *M. persicae* aphids were placed on plants, and one day later adults were removed and 3 1-st instar nymphs were left on the rosette leaves. The total aphid number was counted 14 days after.

Aphid tissue preference test

Thirty alate aphids reared on oil seed rape plants were released inside a cage containing 4 flowering *Arabidopsis* plants of the ecotype Col-0. The aphid release point was approximately 8cm from the cage bottom so that it was positioned at about half the plant height. Twelve days after aphid release, total numbers of aphids were counted in rosette leaves and in stems and flower tissues. This experiment was repeated twice with similar results.

Histochemical localisation VPS52-GUS

Arabidopsis pok mutants were selected as described above and grown under long-day conditions. Flowering plants were placed in individual cages and challenged or unchallenged (control) with aphids for 12 days. Above ground plant tissues were collected and stained with 1mg/ml of 5-bromo-4-chloro-3-indolyl-B-D-glucuronic acid (X-gluc) (# R0852, Thermo Scientific, USA) in X-gluc buffer containing 100mM sodium phosphate buffer pH 7.0, 0.1% Triton X-100, 2mM potassium ferricyanide and 2mM potassium

ferrocyanide. Tissues were vacuum-infiltrated for 10 min and incubated in darkness overnight at 37°C. Chlorophyll was removed by soaking in ethanol. The photographs were taken with a light microscopy. Experiments were repeated three times with similar results.

Quantitative RT-PCR

For the expression analysis of *AtVPS52* in *Arabidopsis*, total RNA was purified using RNeasy Plant mini kit (# 74104, Qiagen, Germany) following manufacturer instructions. cDNA was generated with M-MLV Reverse transcriptase (# M1701, Promega, USA) and samples analyzed by real-time PCR using QuantiTect SYBR[®] Green kit (# 204143, Qiagen, Germany). The primers were designed and analysed with Primer3. Real-time PCR primer sequences were as follows: *AtVPS52_F2*, 5'-AGGAGCCTGCACAAGCTACTTA-3'; *AtVPS52_R2*, 5'-ATGACAGAAAATGGACCCGCA-3'. These primers amplify a fragment of 116 bp. EF1alpha was used as housekeeping gene to calculate relative expression using double delta Ct analysis.

Confocal imaging

Imaging was performed on a Leica TCS-SP2 AOBS (Leica Microsystems) using and HCX PL APO L x40/0.8 water dipping objective and on a Zeiss 710 confocal microscope using a Zeiss PL APO 40x/1.0 water dipping objective. Images were at 1024x1024 resolution and taken using line-by-line sequential scanning. The optimal pinhole diameter and the same gain level for the photomultiplier tube was maintained at all times. ImageJ (NIH) was used for post-acquisition image processing. The excitation wavelength for mRFP was 561nm, its emission was collected from 600 to 630nm. GFP was imaged using 488 nm excitation, and its emission was collected from 500 to 530 nm. CFP was imaged using 405 nm excitation and its emission was collected from 455 to 490 nm. Co-expressed mRFP and GFP as well as co-expressed mRFP and CFP were imaged sequentially using a line-by-line mode.

Y2H reporter assays

Bait-protein encoding vector pDEST32 expressing Mp1 and the prey-encoding vector pDEST22 expressing *AtVPS52* and *SlVPS52* were transformed into the yeast strain MaV203 according to ProQuest[™] Two-Hybrid system protocol (Invitrogen, USA). Transformants were plated onto yeast synthetic drop-out medium (Sigma) lacking Leu and Trp (-LW) and incubated at 30°C for 2 to 3 days. Colonies were picked and cultivated overnight in 5 mL of double drop-out medium (-LW). Dilution series were prepared (10^{-3}) of

each suspension and 5 uL were dropped onto double drop-out medium (-LW), triple drop-out medium lacking Leu, Trp and His (-LWH) plus 3AT (*HIS3* gene inhibitor), and onto a nylon membrane for the X-gal assay along with positive and negative controls according to the ProQuest™ Two-Hybrid system manufacturer's protocol. Plates were incubated at 30°C for 3 to 4 days before photographing. Positive interactions between the expressed proteins resulted in yeast growth and the activation of β -galactosidase in the X-gal membrane assay.

Competing interests

The authors declare that they have no competing interests.

Supplemental Material

Supplemental Figure S1. Protein sequence alignment of StVPS52 with AtVPS52, MtVPS52 and HvVPS52.

Supplemental Figure S2. Protein sequence alignment of Mp1 with Ap1, Mc1 and Rp1.

Supplemental Figure S3. Rp1 and Ap1 do not associate with any VPS52 variant proteins.

Supplemental Figure S4. Limited to no colocalisation of VPS52 with Ara6 and Ara7.

Supplemental Figure S5. Mp1 does not relocalise to vesicle-like compartments upon co-expression with HvVPS52 or MtVPS52.

Supplemental Figure S6. Mp1-like effectors Ap1, Mc1 and Rp1 do not re-localise upon co-expression with any VPS52 variants.

Supplemental Figure S7. Western blot showing expression of GFP-Mp1, GFP-Ap1, GFP-Mc1, RFP-Rp1 and GFP upon agroinfiltration in *Nicotiana benthamiana*.

Supplemental Figure S8. Aphid performance on the *Arabidopsis pok* hemizygous mutant and transcript analyses.

Supplemental Figure S9. Uncropped western blot images corresponding to figures in the main manuscript.

Acknowledgements

We thank P. Birch and E. Huitema for comments and suggestions on the manuscript, M. Armstrong, P. Boevink, S. Engelhardt, P. Hemsley, and K. Wright for technical advice, and T. Warbroek and G. Fraser for technical assistance. We also thank P. Birch for providing the potato Y2H library. We thank S. Bonhomme for kindly providing the Arabidopsis *pok* mutant line.

Figure Legends

Figure 1. *Myzus persicae* effector Mp1 associates with host protein VPS52. (A)

Confirmation of specific two-hybrid interactions in yeast between Mp1 and StVPS52 or AtVPS52 through activation of various reporter genes. Control A represents a negative control, and Control C represents a positive control. -LW represents double drop-out medium lacking Leucine and Tryptophan; -LWH represents triple drop-out medium lacking Leucine, Tryptophan and Histidine; 3AT is the abbreviation of 3-amino-1,2,4-triazole, added to suppress the self-activation of HIS3 gene; X-Gal assay was used to assess the activation of lacZ gene. (B) Co-immunoprecipitation of FLAG-Mp1 and FLAG-Ap1 with GFP-StVPS52 shows that Mp1, but not Ap1, interacts with StVPS52. Leaves of *Nicotiana benthamiana* were infiltrated with *Agrobacterium* strains expressing different combinations of GFP-VPS52 or GFP vector control (GFP-EV) with FLAG-Mp1 or FLAG-Ap1. Three days after infiltration proteins were extracted and subjected to immunoprecipitation with GFP-magnetic beads (IP:GFP) for western blotting with GFP or FLAG antibodies. Lower panel indicates Rubisco stained with Ponceau S (PS) to show equal loading. - indicates absence and + indicates presence of treatment according to upper left panel. This experiment was repeated two times with similar results. The original blots from which images were cropped are shown in Figure S9. (C) Co-immunoprecipitations of FLAG-Mp1 and FLAG-Ap1 with GFP-AtVPS52 shows that Mp1, but not Ap1, interacts with AtVPS52. Leaves of *N. benthamiana* were infiltrated with *Agrobacterium* strains expressing different combinations of GFP-AtVPS52 or GFP vector control (GFP-EV) with FLAG-Mp1 or FLAG-Ap1. Three days after infiltration proteins were extracted and subjected to immunoprecipitation with GFP-magnetic beads (IP:GFP) for western blotting with GFP or FLAG antibodies. Lower panel indicates Rubisco stained with Ponceau S (PS) to show equal loading. - indicates

absence and + indicates presence of treatment according to upper left panel. This experiment was repeated two times with similar results. The original blots from which images were cropped are shown in Figure S9.

Figure 2. *Myzus persicae* effector Mp1 specifically associates with host but not poor-host VPS52s

(A) Co-immunoprecipitations of GFP-StVPS52 with Myc-Mp1, Myc-Ap1, Myc-Mc1 or Myc-Rp1 show that Mp1, but not Mp1-like variants, interacts with StVPS52. Leaves of *Nicotiana benthamiana* were infiltrated with *Agrobacterium* strains expressing different combinations of GFP-StVPS52 with Mp1 variants or GFP vector control (GFP-EV). Three days after infiltration proteins were extracted and subjected to immunoprecipitation with GFP-magnetic beads (IP:GFP) for western blotting with GFP or FLAG antibodies. Lower panel indicates Rubisco stained with Ponceau S (PS) to show equal loading. - indicates absence and + indicates presence of treatment according to upper left panel. The original blots from which images were cropped are shown in Figure S8. (B) Co-immunoprecipitations of FLAG-Mp1 with VPS52 variants St- (*Solanum tuberosum*), At- (*Arabidopsis thaliana*), Hv- (*Hordeum vulgare*) and Mt-StVPS52 (*Medicago truncatula*) show that Mp1 interacts only with StVPS52 and AtVPS52. Leaves of *N. benthamiana* were infiltrated with *Agrobacterium* strains expressing different combinations of GFP-StVPS52 with FLAG-Mp1 or GFP vector control (GFP-EV). Three days after infiltration proteins were extracted and subjected to immunoprecipitation with GFP-magnetic beads (IP:GFP) for western blotting with GFP or FLAG antibodies. Lower panel indicates Rubisco stained with Ponceau S (PS) to show equal loading. - indicates absence and + indicates presence of treatment according to upper left panel. The original blots from which images were cropped are shown in Figure S9.

Figure 3. Overexpression of host VPS52s causes relocalization of Mp1 to vesicle-like structures that associate with the prevacuolar compartment

(A) Confocal microscopy images of *Nicotiana benthamiana* leaves transiently expressing GFP-AtVPS52 in combination with subcellular marker CFP-PS1. Images were taken 3 days after agroinfiltration. Merged figures represent the overlay images of the GFP with CFP channels. Spatial plot profiles represent the colocalisation level across a defined region of interest (ROI) depicted in the merged images. Plot profiles were done with Fiji software (NHI). (B) Confocal microscopy images of *N. benthamiana* leaves transiently expressing GFP-StVPS52 in combination with subcellular marker CFP-PS1. Images were taken 3

days after agroinfiltration Merged figures represent the overlay images of the GFP with CFP channels. Spatial plot profiles represent the colocalisation level across a defined region of interest (ROI) depicted in the merged images. Plot profiles were done with Fiji software (NHI). (C) Colocalization of GFP-Mp1 with mRFP-StVPS52, mRFP-AtVPS52, or mRFP by confocal microscopy. Images were taken 3 days after agroinfiltration of *N. benthamiana* leaves. Merged figures represent the overlay images of the GFP and mRFP channels. Scale bars= 20 μ m

Figure 4. *Myzus persicae* effector Mp1, but not Mp1-like effectors from other aphid species, promotes virulence on *Nicotiana benthamiana* upon phloem-specific overexpression

Leaves of *Nicotiana benthamiana* transiently expressing different Mp1-like effectors under the phloem-specific AtSUC2 promoter were challenged with *M. persicae*. Aphid fecundity was assessed over a 14-day period. Ap1, Rp1, and Mc1, are the Mp1-like effectors from *Acyrtosiphon pisum*, *Rhopalosiphum padi*, and *Myzus cerasi*, respectively. Empty vector (EV) was used as a control. The graph shows the means of 4 independent replicates, where $n=12$ represent the maximum number of samples per treatment in each replicate. The normally distributed data set was treated with One-way ANOVA Welch and Brown-Forsythe tests for unequal variances, and Scheffe Post-Hoc test for unequal group size. Error bars represent the standard error of the means and asterisks (***) indicates significant differences between treatments versus vector control ($P<0.01$).

Figure 5. StVPS52 reduce *Myzus persicae* virulence on *Nicotiana benthamiana* upon phloem-specific overexpression

Leaves of *N. benthamiana* transiently expressing StVPS52 (potato) under the phloem-specific AtSUC2 promoter were challenged with *M. persicae*. Aphid fecundity was assessed over a 14-day period. Empty vector (EV) was used as a control. Graphs represent the means of 3 independent replicates, where $n=12$ represents the number of samples per treatment in each replicate, and error bars represent the standard error of the means. Asterisks (*) indicate significant differences between treatment and the control (t test, $P<0.05$).

Figure 6. Aphid infestation causes degradation of VPS52. (A) *pok* mutant plants were infested with *Myzus persicae* for 5-7 days and collected for GUS-staining. Images were taken with a light microscopy. The experiment was repeated three times with similar

results. (B) *N. benthamiana* leaves transiently over-expressing GFP vector control (GFP-EV), GFP-StVPS52, GFP-AtVPS52 in combination with FLAG-Mp1 or infested with *M. persicae* for 3 days were collected for protein extraction and western blotting with anti-GFP and anti-FLAG antibodies. Lower panel indicates Rubisco stained with Ponceau S (PS) to show equal loading. - indicates absence and + indicates presence of treatment according to upper left panel. (C) *N. benthamiana* leaves transiently over-expressing GFP vector control (GFP-EV) and GFP-StVPS52 were challenged with *Phytophthora capsici* zoospores prior to protein extraction and western blotting with GFP antibodies. *N. benthamiana* leaves transiently over-expressing GFP vector control (GFP-EV) and GFP-StVPS52 were challenged with the PAMP flg22 prior to protein extraction and western blotting with GFP antibodies. Lower panel indicates Rubisco stained with Ponceau S (PS) to show equal loading. - indicates absence and + indicates presence of treatment according to upper left panel.

Figure 7. Aphids preferentially colonize Arabidopsis stems and flower tissues over rosette leaves. Aphid numbers on rosette leaves or stem and flower tissues of flowering plants 12 days after the release of alate adults. Graph represents the means of 2 independent replicates, where $n=4$ represents the number of plants in each replicate, and error bars represent the standard error of the means. Asterisks (*) indicate significant differences between rosette leaves and stems plus flowers tissues (paired t test, $P<0.05$).

References

- Ashouri A, Michaud D, Cloutier C (2001) Unexpected effects of different potato resistance factors to the Colorado Potato Beetle (Coleoptera: Chrysomelidae) on the Potato Aphid (Homoptera: Aphididae). *Environmental Entomology* 30: 524-532
- Atamian HS, Chaudhary R, Dal Cin V, Bao E, Girke T, Kaloshian I (2013) In planta expression or delivery of potato Aphid macrosiphum euphorbiae effectors Me10 and Me23 enhances aphid fecundity. *MPMI* 26: 67-74
- Ben Khaled S, Postma J, Robatzek S (2015) A moving view: Subcellular trafficking processes in pattern recognition receptor-triggered plant immunity. *Annu Rev Phytopathol* 53: 379-402
- Boevink PC, Wang X, McLellan H, He Q, Naqvi S, Armstrong MR, Zhang W, Hein I, Gilroy EM, Tian Z, Birch PR (2016) A *Phytophthora infestans* RXLR effector targets plant PP1c isoforms that promote late blight disease. *Nat Commun* 7: 10311
- Bos JI, Armstrong MR, Gilroy EM, Boevink PC, Hein I, Taylor RM, Zhendong T, Engelhardt S, Vetukuri RR, Harrower B, Dixelius C, Bryan G, Sadanandom A, Whisson SC, Kamoun S, Birch PR (2010) *Phytophthora infestans* effector AVR3a is essential for virulence and manipulates plant immunity by stabilizing host E3 ligase CMPG1. *Proc Natl Acad Sci U S A* 107: 9909-9914

720 Bos JIB, Prince DC, Pitino M, Maffei mE, Win J, Hogenhout SA (2010) A functional
 721 genomics approach identifies candidate effectors from the aphid species *Myzus*
 722 *persicae* (Green Peach Aphid). *PLoS Genetics* 6(11): e1001216
 723 Bozkurt TO, Schornack S, Win J, Shindo T, Ilyas M, Oliva R, Cano LM, Jones AM,
 724 Huitema E, van der Hoorn RA, Kamoun S (2011) *Phytophthora infestans* effector
 725 AVRblb2 prevents secretion of a plant immune protease at the haustorial interface.
 726 *Proc Natl Acad Sci U S A* 108: 20832-20837
 727 Carolan J, Caragea D, Reardon KT, Mutti NS, Dittmer N, Pappan K, Cui F, Castaneto M,
 728 Poulain J, Dossat C, Tagu D, Reese J, Reeck G, Wilkinson TL, Edwards OR (2011)
 729 Predicted effector molecules in the salivary secretome of the Pea Aphid
 730 (*Acyrtosiphon pisum*): A dual transcriptomic/proteomic approach. *Journal of*
 731 *Proteome Research* 10: 1505-1518
 732 Carolan J, Fitzroy CIJ, Ashton PD, Douglas AE, Wilkinson TL (2009) The secreted salivary
 733 proteome of the pea aphid *Acyrtosiphon pisum* characterised by mass
 734 spectrometry. *Proteomics* 9: 2457-2467
 735 Conibear E, Cleck J, Stevens TH (2003) Vps51p mediates the association of the GARP
 736 (Vps52/53/54) complex with the late Golgi t-SNARE Tlg1p. *Mol Biol Cell* 14: 1610-
 737 1623
 738 Conibear E, Stevens TH (2000) Vps52p, Vps53p, and Vps54p form a novel multisubunit
 739 complex required for protein sorting at the yeast late Golgi. *Mol Biol Cell* 11: 305-
 740 323
 741 Dagdas YF, Belhaj K, Maqbool A, Chaparro-Garcia A, Pandey P, Petre B, Tabassum N,
 742 Cruz-Mireles N, Hughes RK, Sklenar J, Win J, Menke F, Findlay K, Banfield MJ,
 743 Kamoun S, Bozkurt TO (2016) An effector of the Irish potato famine pathogen
 744 antagonizes a host autophagy cargo receptor. *Elife* 5:e10856
 745 Davis JA, Radcliffe EB (2008) Reproduction and feeding behavior of *Myzus persicae* on
 746 four cereals. *J Econ Entomol* 101: 9-16
 747 Elzinga DA, De Vos M, Jander G (2014) Suppression of plant defenses by a *Myzus*
 748 *persicae* (Green Peach Aphid) salivary effector protein. *MPMI* 27: 747-756
 749 Elzinga DA, Jander G (2013) The role of protein effectors in plant-aphid interactions
 750 *Curr Opin Plant Biol* 16:451-456.
 751 Frohlich A, Gaupels F, Sarioglu H, Holzmeister C, Spannagl M, Durner J, Lindermayr C
 752 (2012) Looking deep inside: detection of low-abundance proteins in leaf extracts of
 753 *Arabidopsis* and phloem exudates of pumpkin. *Plant Physiol* 159: 902-914
 754 Gao LL, Horbury R, Nair RM, Singh KB, Edwards OR (2007) Characterization of
 755 resistance to multiple aphid species (Hemiptera: Aphididae) in *Medicago truncatula*.
 756 *Bull Entomol Res* 97: 41-48
 757 Girousse C, Moulia B, Silk W, Bonnemain J (2005) Aphid infestation causes different
 758 changes in carbon and nitrogen allocation in alfalfa stems as well as different
 759 inhibitions of longitudinal and radial expansion1. *Plant Physiol*. 137: 1474-1484
 760 Gottwald JR, Krysan PJ, Young JC, Evert RF, Sussman MR (2000) Genetic evidence for
 761 the in planta role of phloem-specific plasma membrane sucrose transporters. *Proc*
 762 *Natl Acad Sci U S A* 97: 13979-13984
 763 Groll M, Schellenberg B, Bachmann AS, Archer CR, Huber R, Powell TK, Lindow S, Kaiser
 764 M, Dudler R (2008) A plant pathogen virulence factor inhibits the eukaryotic
 765 proteasome by a novel mechanism. *Nature* 452: 755-758
 766 Gu Y, Innes RW (2012) The KEEP ON GOING protein of *Arabidopsis* regulates
 767 intracellular protein trafficking and is degraded during fungal infection. *Plant Cell* 24:
 768 4717-4730
 769 Guermontprez H, Smertenko A, Crosnier M, Durandet M, Vrielynck N, Guerche P, Hussey
 770 P, Satiat-Jeunemaitre B, Bonhomme S (2008) The POK/AtVPS52 protein localizes

771 to several distinct post-Golgi compartments in sporophytic gametophytic cells. *J.*
772 *Exp Bot.* 59: 3087-3098

773 Guldemon JA, van der Brink WJ, den Belder E (1998) Methods of assessing population
774 increase in aphids and the effect of growthstage of the host plant on population
775 growth rates. *Entomologia Experimentalis et Applicata* 86: 163-173

776 Harmel N, Létocart E, Cherqui A, Giordanengo P, Mazzucchelli G, Guillonneau F, De
777 Pauw E, Haubruge E, Francis F (2008) Identification of aphid salivary proteins: a
778 proteomic investigation of *Myzus persicae*. *Insect Molec Biol.* 17: 165-174

779 Heard W, Sklenář J, Tomé D, Robatzek S, Jones AM (2015) Identification of regulatory
780 and cargo proteins of endosomal and secretory pathways in *Arabidopsis thaliana* by
781 proteomic dissection. *Mol Cell Proteomics* 14: 1796-1813

782 Imlau A, Truernit E, Sauer N (1999) Cell-to-cell and long-distance trafficking of the green
783 fluorescent protein in the phloem and symplastic unloading of the protein into sink
784 tissues. *Plant Cell* 11: 309-322

785 Jiang S, Yao J, Ma KW, Zhou H, Song J, He SY, Ma W (2013) Bacterial effector activates
786 jasmonate signaling by directly targeting JAZ transcriptional repressors. *PLoS*
787 *Pathog* 9: e1003715

788 Karimi M, Inze D, Depicker A (2002) GATEWAY vectors for *Agrobacterium*-mediated plant
789 transformation. *Trends in Plant Science* 7: 193-195

790 King SR, McLellan H, Boevink PC, Armstrong MR, Bukharova T, Sukarta O, Win J,
791 Kamoun S, Birch PR, Banfield MJ (2014) *Phytophthora infestans* RXLR effector
792 PexRD2 interacts with host MAPKKK epsilon to suppress plant immune signaling.
793 *Plant Cell* 26: 1345-1359

794 Kotzer AM, Brandizzi F, Neumann U, Paris N, Moore I, Hawes C (2004) AtRabF2b (Ara7)
795 acts on the vacuolar trafficking pathway in tobacco leaf epidermal cells. *J Cell Sci*
796 117: 6377-6389

797 Lee AH, Hurley B, Felsensteiner C, Yea C, Ckurshumova W, Bartetzko V, Wang PW,
798 Quach V, Lewis JD, Liu YC, Bornke F, Angers S, Wilde A, Guttman DS, Desveaux
799 D (2012) A bacterial acetyltransferase destroys plant microtubule networks and
800 blocks secretion. *PLoS Pathog* 8: e1002523

801 Lin MK, Lee YJ, Lough TJ, Phinney BS, Lucas WJ (2009) Analysis of the pumpkin phloem
802 proteome provides insights into angiosperm sieve tube function. *Mol Cell*
803 *Proteomics* 8: 343-356

804 Lobstein E, Guyon A, Férault M, Twell D, Pelletier G, Bonhomme S (2004) The putative
805 *Arabidopsis* homolog of yeast vps52p is required for pollen tube elongation,
806 localizes to Golgi, and might be involved in vesicle trafficking. *Plant Physiol* 135:
807 1480-1490

808 Mutti NS, Louis J, Pappan K, Begum K, Chen M-S, Park Y, Dittmer N, Marshall J, Reese
809 J, Reeck G (2008) A protein from the salivary glands of the pea aphid,
810 *Acyrtosiphon pisum*, is essential in feeding on a host plant. *Proc Natl Acad Sci U S*
811 *A.* 105: 9965-9969

812 Nakagawa T, Kurose T, Hino T, Tanaka K, Kawamukai M, Niwa Y, Toyooka K, Matsuoka
813 K, Jinbo T, Kimura T (2007) Development of series of gateway binary vectors,
814 pGWBs, for realizing efficient construction of fusion genes for plant transformation.
815 *J Biosci Bioeng.* 104: 34-41

816 Nomura K, Debroy S, Lee YH, Pumplin N, Jones J, He SY (2006) A bacterial virulence
817 protein suppresses host innate immunity to cause plant disease. *Science* 313: 220-
818 223

819 Pahari S, Cormark RD, Blackshaw MT, Liu C, Erickson JL, Schultz EA (2014) *Arabidopsis*
820 UNHINGED encodes a VPS51 homolog and reveals a role for the GARP complex
821 in leaf shape and vein patterning. *Development* 141: 1894-1905

- Pitino M, Hogenhout SA (2013) Aphid protein effectors promote aphid colonization in a plant species-specific manner. *MPMI* 26: 130-139
- Plett JM, Daguerre Y, Wittulsky S, Vayssieres A, Deveau A, Melton SJ, Kohler A, Morrell-Falvey JL, Brun A, Veneault-Fourrey C, Martin F (2014) Effector MiSSP7 of the mutualistic fungus *Laccaria bicolor* stabilizes the *Populus* JAZ6 protein and represses jasmonic acid (JA) responsive genes. *Proc Natl Acad Sci U S A* 111: 8299-8304
- Reggiori F, Wang C, Stromhaug P, Shintani T, Klionsky D (2003) Vps51 is part of the yeast Vps fifty-three tethering complex essential for retrograde traffic from the early endosome and Cvt vesicle completion. *J Biol Chem* 278: 5009-5020
- Rodriguez PA, Bos JI (2013) Toward understanding the role of aphid effectors in plant infestation. *Mol Plant Microbe Interact* 26: 25-30
- Rodriguez PA, Stam R, Warbroek T, Bos JIB (2014) Mp10 and Mp42 from the aphid species *Myzus persicae* trigger plant defenses in *Nicotiana benthamiana* through different activities. *MPMI* 27: 30-39
- Sandström J, Telang A, Moran NA (2000) Nutritional enhancement of host plants by aphids - a comparison of three aphid species on grasses. *J Insect Physiol* 46: 33-40
- Schindler C, Chen Y, Pu J, Guo X, Bonifacino JS (2015) EARP is a multisubunit tethering complex involved in endocytic recycling. *Nat Cell Biol* 17: 639-650
- Schulze-Lefert P, Panstruga R (2011) A molecular evolutionary concept connecting nonhost resistance, pathogen host range, and pathogen speciation. *Trends Plant Sci* 16: 117-125
- Song J, Win J, Tian M, Schornack S, Kaschani F, Ilyas M, van der Hoorn RA, Kamoun S (2009) Apoplastic effectors secreted by two unrelated eukaryotic plant pathogens target the tomato defense protease Rcr3. *Proc Natl Acad Sci U S A* 106: 1654-1659
- Teh OK, Hofius D (2014) Membrane trafficking and autophagy in pathogen-triggered cell death and immunity. *J Exp Bot* 65: 1297-1312
- Thorpe P, Cock PJ, Bos J (2016) Comparative transcriptomics and proteomics of three different aphid species identifies core and diverse effector sets. *BMC Genomics* 17: 172
- Ustun S, Bartetzko V, Bornke F (2013) The *Xanthomonas campestris* type III effector XopJ targets the host cell proteasome to suppress salicylic-acid mediated plant defence. *PLoS Pathog* 9: e1003427
- Zheng X, McLellan H, Fraiture M, Liu X, Boevink PC, Gilroy EM, Chen Y, Kandel K, Sessa G, Birch PR, Brunner F (2014) Functionally redundant RXLR effectors from *Phytophthora infestans* act at different steps to suppress early flg22-triggered immunity. *PLoS Pathog* 10: e1004057




Figure 1. *Myzus persicae* effector Mp1 associates with host protein VPS52. (A) Confirmation of specific two-hybrid interactions in yeast between Mp1 and StVPS52 or AtVPS52 through activation of various reporter genes. Control A represents a negative control, and Control C represents a positive control. -LW represents double drop-out medium lacking Leucine and Tryptophan; -LWH represents triple drop-out medium lacking Leucine, Tryptophan and Histidine; 3AT is the abbreviation of 3-amino-1,2,4-triazole, added to suppress the self-activation of *HIS3* gene; X-Gal assay was used to assess the activation of *lacZ* gene. (B) Co-immunoprecipitation of FLAG-Mp1 and FLAG-Ap1 with GFP-StVPS52 shows that Mp1, but not Ap1, interacts with StVPS52. Leaves of *Nicotiana benthamiana* were infiltrated with *Agrobacterium* strains expressing different combinations of GFP-VPS52 or GFP vector control (GFP-EV) with FLAG-Mp1 or FLAG-Ap1. Three days after infiltration proteins were extracted and subjected to immunoprecipitation with GFP-magnetic beads (IP:GFP) for western blotting with GFP or FLAG antibodies. Lower panel indicates Rubisco stained with Ponceau S (PS) to show equal loading. - indicates absence and + indicates presence of treatment according to upper left panel. This experiment was repeated two times with similar results. The original blots from which images were cropped are shown in Figure S8. (C) Co-immunoprecipitations of FLAG-Mp1 and FLAG-Ap1 with GFP-AtVPS52 shows that Mp1, but not Ap1, interacts with AtVPS52. Leaves of *N. benthamiana* were infiltrated with *Agrobacterium* strains expressing different combinations of GFP-AtVPS52 or GFP vector control (GFP-EV) with FLAG-Mp1 or FLAG-Ap1. Three days after infiltration proteins were extracted and subjected to immunoprecipitation with GFP-magnetic beads (IP:GFP) for western blotting with GFP or FLAG antibodies. Lower panel indicates Rubisco stained with Ponceau S (PS) to show equal loading. - indicates absence and + indicates presence of treatment according to upper left panel. This experiment was repeated two times with similar results. The original blots from which images were cropped are shown in Figure S9




Figure 2. *Myzus persicae* effector Mp1 specifically associates with host but not poor-host VPS52s (A) Co-immunoprecipitations of GFP-StVPS52 with Myc-Mp1, Myc-Ap1, Myc-Mc1 or Myc-Rp1 show that Mp1, but not Mp1-like variants, interacts with StVPS52 Leaves of *Nicotiana benthamiana* were infiltrated with *Agrobacterium* strains expressing different combinations of GFP-StVPS52 with Mp1 variants or GFP vector control (GFP-EV). Three days after infiltration proteins were extracted and subjected to immunoprecipitation with GFP-magnetic beads (IP:GFP) for western blotting with GFP or FLAG antibodies. Lower panel indicates Rubisco stained with Ponceau S (PS) to show equal loading. - indicates absence and + indicates presence of treatment according to upper left panel. The original blots from which images were cropped are shown in Figure S8. (B) Co-immunoprecipitations of FLAG-Mp1 with VPS52 variants St- (potato), At- (*Arabidopsis thaliana*), Hv- (*Hordeum vulgare*) and Mt-StVPS52 (*Medicago truncatula*) show that Mp1 interacts only with StVPS52 and AtVPS52. Leaves of *Nicotiana benthamiana* were infiltrated with *Agrobacterium* strains expressing different combinations of GFP-StVPS52 with FLAG-Mp1 or GFP vector control (GFP-EV). Three days after infiltration proteins were extracted and subjected to immunoprecipitation with GFP-magnetic beads (IP:GFP) for western blotting with GFP or FLAG antibodies. Lower panel indicates Rubisco stained with Ponceau S (PS) to show equal loading. - indicates absence and + indicates presence of treatment according to upper left panel. The original blots from which images were cropped are shown in Figure S9.




Figure 3. Overexpression of host VPS52s causes relocation of Mp1 to vesicle-like structures that associate with the prevacuolar compartment (A) Confocal microscopy images of *Nicotiana benthamiana* leaves transiently expressing GFP-AtVPS52 in combination with subcellular marker CFP-PS1. Images were taken 3 days after agroinfiltration. Merged figures represent the overlay images of the GFP with CFP channels. Spatial plot profiles represent the colocalisation level across a defined region of interest (ROI) depicted in the merged images. Plot profiles were done with Fiji software (NHI). (B) Confocal microscopy images of *N. benthamiana* leaves transiently expressing GFP-StVPS52 in combination with subcellular marker CFP-PS1. Images were taken 3 days after agroinfiltration. Merged figures represent the overlay images of the GFP with CFP channels. Spatial plot profiles represent the colocalisation level across a defined region of interest (ROI) depicted in the merged images. Plot profiles were done with Fiji software (NHI). (C) Colocalization of GFP-Mp1 with mRFP-StVPS52, mRFP-AtVPS52, or mRFP by confocal microscopy. Images were taken 3 days after agroinfiltration of *Nicotiana benthamiana* leaves. Merged figures represent the overlay images of the GFP and mRFP channels. Scale bars= 20 μ m




Figure 4. *Myzus persicae* effector Mp1, but not Mp1-like effectors from other aphid species, promotes virulence on *Nicotiana benthamiana* upon phloem-specific overexpression.

Leaves of *Nicotiana benthamiana* transiently expressing different Mp1-like effectors under the phloem-specific AtSUC2 promoter were challenged with *M. persicae*. Aphid fecundity was assessed over a 14-day period. Ap1, Rp1, and Mc1, are the Mp1-like effectors from *Acyrtosiphon pisum*, *Rhopalosiphum padi*, and *Myzus cerasi*, respectively. Empty vector (EV) was used as a control. The graph shows the means of 4 independent replicates, where n=12 represent the maximum number of samples per treatment in each replicate.

The normally distributed data set was treated with One-way ANOVA Welch and Brown-Forsythe tests for unequal variances, and Scheffe Post-Hoc test for unequal group size. Error bars represent the standard error of the means and asterisks (***) indicates significant differences between treatments and the vector control ($P < 0.01$).





Figure 5. StVPS52 reduce *Myzus persicae* virulence on *Nicotiana benthamiana* upon phloem-specific overexpression.

Leaves of *N. benthamiana* transiently expressing StVPS52 (potato) under the phloem-specific AtSUC2 promoter were challenged with *M. persicae*. Aphid fecundity was assessed over a 14-day period. Empty vector (EV) was used as a control. Graphs represent the means of 3 independent replicates, where n=12 represents the number of samples per treatment in each replicate, and error bars represent the standard error of the means. Asterisks (*) indicate significant differences between treatment and the control (t test, $P < 0.05$).

AAtVPS52::GUS - *M. persicae*AtVPS52::GUS + *M. persicae***B**

| | | | | | | | | | |
|--------------------|---|---|---|---|---|---|---|---|---|
| GFP-EV | - | - | - | - | - | - | + | + | + |
| GFP-StVPS52 | + | - | + | - | + | - | - | - | - |
| GFP-AtVPS52 | - | + | - | + | - | + | - | - | - |
| FLAG-Mp1 | - | - | + | + | - | - | - | + | - |
| <i>M. persicae</i> | - | - | - | - | + | + | - | - | + |

**C**

| | | | | |
|-------------------|---|---|---|---|
| GFP-StVPS52 | + | + | - | - |
| GFP-EV | - | - | + | + |
| <i>P. capsici</i> | - | + | - | + |

| | | | | |
|-------------|---|---|---|---|
| GFP-StVPS52 | + | + | - | - |
| GFP-EV | - | - | + | + |
| flg22 | - | + | - | + |




Figure 6. Aphid infestation causes degradation of VPS52. (A) pok mutant plants were infested with *Myzus persicae* for 5-7 days and collected for GUS-staining. Images were taken with a light microscopy. The experiment was repeated three times with similar results. (B) *Nicotiana benthamiana* leaves transiently over-expressing GFP vector control (GFP-EV), GFP-StVPS52, GFP-AtVPS52 in combination with FLAG-Mp1 or infested with *M. persicae* for 3 days were collected for protein extraction and western blotting with anti-GFP and anti-FLAG antibodies. Lower panel indicates Rubisco stained with Ponceau S (PS) to show equal loading. - indicates absence and + indicates presence of treatment according to upper left panel. (C) *N. benthamiana* leaves transiently over-expressing GFP vector control (GFP-EV) and GFP-StVPS52 were challenged with *Phytophthora capsici* zoospores prior to protein extraction and western blotting with GFP antibodies. *N. benthamiana* leaves transiently over-expressing GFP vector control (GFP-EV) and GFP-StVPS52 were challenged with the PAMP flg22 prior to protein extraction and western blotting with GFP antibodies. Lower panel indicates Rubisco stained with Ponceau S (PS) to show equal loading. - indicates absence and + indicates presence of treatment according to upper left panel.



Figure 7. Aphids preferentially colonize Arabidopsis stems and flower tissues over rosette leaves. Aphid numbers on rosette leaves or stem and flower tissues of flowering plants 12 days after the release of alate adults. Graph represents the means of 2 independent replicates, where $n=4$ represents the number of plants in each replicate, and error bars represent the standard error of the means. Asterisks (*) indicate significant differences between rosette leaves and stems plus flowers tissues (paired t test, $P<0.05$).

Parsed Citations

Ashouri A, Michaud D, Cloutier C (2001) Unexpected effects of different potato resistance factors to the Colorado Potato Beetle (Coleoptera: Chrysomelidae) on the Potato Aphid (Homoptera: Aphididae). Environmental Entomology 30: 524-532

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Atamian HS, Chaudhary R, Dal Cin V, Bao E, Girke T, Kaloshian I (2013) In planta expression or delivery of potato Aphid *macrosiphum euphorbiae* effectors Me10 and Me23 enhances aphid fecundity. MPMI 26: 67-74

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Ben Khaled S, Postma J, Robatzek S (2015) A moving view: Subcellular trafficking processes in pattern recognition receptor-triggered plant immunity. Annu Rev Phytopathol 53: 379-402

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Boevink PC, Wang X, McLellan H, He Q, Naqvi S, Armstrong MR, Zhang W, Hein I, Gilroy EM, Tian Z, Birch PR (2016) A *Phytophthora infestans* RXLR effector targets plant PP1c isoforms that promote late blight disease. Nat Commun 7: 10311

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Bos JI, Armstrong MR, Gilroy EM, Boevink PC, Hein I, Taylor RM, Zhendong T, Engelhardt S, Vetukuri RR, Harrower B, Dixelius C, Bryan G, Sadanandom A, Whisson SC, Kamoun S, Birch PR (2010) *Phytophthora infestans* effector AVR3a is essential for virulence and manipulates plant immunity by stabilizing host E3 ligase CMPG1. Proc Natl Acad Sci U S A 107: 9909-9914

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Bos JIB, Prince DC, Pitino M, Maffei mE, Win J, Hogenhout SA (2010) A functional genomics approach identifies candidate effectors from the aphid species *Myzus persicae* (Green Peach Aphid). PLOs Genetics 6(11): e1001216

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Bozkurt TO, Schornack S, Win J, Shindo T, Ilyas M, Oliva R, Cano LM, Jones AM, Huitema E, van der Hoorn RA, Kamoun S (2011) *Phytophthora infestans* effector AVRblb2 prevents secretion of a plant immune protease at the haustorial interface. Proc Natl Acad Sci U S A 108: 20832-20837

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Carolan J, Caragea D, Reardon KT, Mutti NS, Dittmer N, Pappan K, Cui F, Castaneto M, Poulain J, Dossat C, Tagu D, Reese J, Reeck G, Wilkinson TL, Edwards OR (2011) Predicted effector molecules in the salivary secretome of the Pea Aphid (*Acyrtosiphon pisum*): A dual transcriptomic/proteomic approach. Journal of Proteome Research 10: 1505-1518

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Carolan J, Fitzroy CIJ, Ashton PD, Douglas AE, Wilkinson TL (2009) The secreted salivary proteome of the pea aphid *Acyrtosiphon pisum* characterised by mass spectrometry. Proteomics 9: 2457-2467

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Conibear E, Cleck J, Stevens TH (2003) Vps51p mediates the association of the GARP (Vps52/53/54) complex with the late Golgi t-SNARE Tlg1p. Mol Biol Cell 14: 1610-1623

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Conibear E, Stevens TH (2000) Vps52p, Vps53p, and Vps54p form a novel multisubunit complex required for protein sorting at the yeast late Golgi. Mol Biol Cell 11: 305-323

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Dagdas YF, Belhaj K, Maqbool A, Chaparro-Garcia A, Pandey P, Petre B, Tabassum N, Cruz-Mireles N, Hughes RK, Sklenar J, Win J, Menke F, Findlay K, Banfield MJ, Kamoun S, Bozkurt TO (2016) An effector of the Irish potato famine pathogen antagonizes a host autophagy cargo receptor. Elife 5:e10856

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Davis JA, Radcliffe EB (2008) Reproduction and feeding behavior of *Myzus persicae* on four cereals. J Econ Entomol 101: 9-16

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Elzinga DA, De Vos M, Jander G (2014) Suppression of plant defenses by a Myzus persicae (Green Peach Aphid) salivary effector protein. MPMI 27: 747-756

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Elzinga DA, Jander G (2013) The role of protein effectors in plant-aphid interactions

Curr Opin Plant Biol 16:451-456.

Frohlich A, Gaupels F, Sarioglu H, Holzmeister C, Spannagl M, Durner J, Lindermayr C (2012) Looking deep inside: detection of low-abundance proteins in leaf extracts of Arabidopsis and phloem exudates of pumpkin. Plant Physiol 159: 902-914

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Gao LL, Horbury R, Nair RM, Singh KB, Edwards OR (2007) Characterization of resistance to multiple aphid species (Hemiptera: Aphididae) in Medicago truncatula. Bull Entomol Res 97: 41-48

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Girousse C, Moulia B, Silk W, Bonnemain J (2005) Aphid infestation causes different changes in carbon and nitrogen allocation in alfalfa stems as well as different inhibitions of longitudinal and radial expansion1. Plant Physiol. 137: 1474-1484

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Gottwald JR, Krysan PJ, Young JC, Evert RF, Sussman MR (2000) Genetic evidence for the in planta role of phloem-specific plasma membrane sucrose transporters. Proc Natl Acad Sci U S A 97: 13979-13984

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Groll M, Schellenberg B, Bachmann AS, Archer CR, Huber R, Powell TK, Lindow S, Kaiser M, Dudler R (2008) A plant pathogen virulence factor inhibits the eukaryotic proteasome by a novel mechanism. Nature 452: 755-758

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Gu Y, Innes RW (2012) The KEEP ON GOING protein of Arabidopsis regulates intracellular protein trafficking and is degraded during fungal infection. Plant Cell 24: 4717-4730

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Guermonprez H, Smertenko A, Crosnier M, Durandet M, Vrielynck N, Guerche P, Hussey P, Satiat-Jeunemaitre B, Bonhomme S (2008) The POK/AtVPS52 protein localizes to several distinct post-Golgi compartments in sporophytic gametophytic cells. J. Exp Bot. 59: 3087-3098

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Guldmond JA, van der Brink WJ, den Belder E (1998) Methods of assessing population increase in aphids and the effect of growthstage of the host plant on population growth rates. Entomologia Experimentalis et Applicata 86: 163-173

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Harmeï N, Létocart E, Cherqui A, Giordanengo P, Mazzucchelli G, Guillonéau F, De Pauw E, Haubruge E, Francis F (2008) Identification of aphid salivary proteins: a proteomic investigation of Myzus persicae. Insect Molec Biol. 17: 165-174

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Heard W, Sklenár J, Tomé D, Robatzek S, Jones AM (2015) Identification of regulatory and cargo proteins of endosomal and secretory pathways in Arabidopsis thaliana by proteomic dissection. Mol Cell Proteomics 14: 1796-1813

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Imlau A, Truernit E, Sauer N (1999) Cell-to-cell and long-distance trafficking of the green fluorescent protein in the phloem and symplastic unloading of the protein into sink tissues. Plant Cell 11: 309-322

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Jiang S, Yao J, Ma KW, Zhou H, Song J, He SY, Ma W (2013) Bacterial effector activates jasmonate signaling by directly targeting JAZ transcriptional repressors. PLoS Pathog 9: e1003715

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Karimi M, Inze D, Depicker A (2002) GATEWAY vectors for Agrobacterium-mediated plant transformation. Trends in Plant Science 7: 193-195

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

King SR, McLellan H, Boevink PC, Armstrong MR, Bukharova T, Sukarta O, Win J, Kamoun S, Birch PR, Banfield MJ (2014) Phytophthora infestans RXLR effector PexRD2 interacts with host MAPKKK epsilon to suppress plant immune signaling. Plant Cell 26: 1345-1359

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Kotzer AM, Brandizzi F, Neumann U, Paris N, Moore I, Hawes C (2004) AtRabF2b (Ara7) acts on the vacuolar trafficking pathway in tobacco leaf epidermal cells. J Cell Sci 117: 6377-6389

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Lee AH, Hurley B, Felsensteiner C, Yea C, Ckurshumova W, Bartetzko V, Wang PW, Quach V, Lewis JD, Liu YC, Bornke F, Angers S, Wilde A, Guttman DS, Desveaux D (2012) A bacterial acetyltransferase destroys plant microtubule networks and blocks secretion. PLoS Pathog 8: e1002523

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Lin MK, Lee YJ, Lough TJ, Phinney BS, Lucas WJ (2009) Analysis of the pumpkin phloem proteome provides insights into angiosperm sieve tube function. Mol Cell Proteomics 8: 343-356

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Lobstein E, Guyon A, Féralut M, Twell D, Pelletier G, Bonhomme S (2004) The putative Arabidopsis homolog of yeast vps52p is required for pollen tube elongation, localizes to Golgi, and might be involved in vesicle trafficking. Plant Physiol 135: 1480-1490

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Mutti NS, Louis J, Pappan K, Begum K, Chen M-S, Park Y, Dittmer N, Marshall J, Reese J, Reeck G (2008) A protein from the salivary glands of the pea aphid, Acyrthosiphon pisum, is essential in feeding on a host plant. Proc Natl Acad Sci U S A 105: 9965-9969

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Nakagawa T, Kurose T, Hino T, Tanaka K, Kawamukai M, Niwa Y, Toyooka K, Matsuoka K, Jinbo T, Kimura T (2007) Development of series of gateway binary vectors, pGWBs, for realizing efficient construction of fusion genes for plant transformation. J Biosci Bioeng. 104: 34-41

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Nomura K, Debroy S, Lee YH, Pumlun N, Jones J, He SY (2006) A bacterial virulence protein suppresses host innate immunity to cause plant disease. Science 313: 220-223

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Pahari S, Cormark RD, Blackshaw MT, Liu C, Erickson JL, Schultz EA (2014) Arabidopsis UNHINGED encodes a VPS51 homolog and reveals a role for the GARP complex in leaf shape and vein patterning. Development 141: 1894-1905

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Pitino M, Hogenhout SA (2013) Aphid protein effectors promote aphid colonization in a plant species-specific manner. MPMI 26: 130-139

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Plett JM, Daguerre Y, Wittulsky S, Vayssieres A, Deveau A, Melton SJ, Kohler A, Morrell-Falvey JL, Brun A, Veneault-Fourrey C, Martin F (2014) Effector MiSSP7 of the mutualistic fungus Laccaria bicolor stabilizes the Populus JAZ6 protein and represses jasmonic acid (JA) responsive genes. Proc Natl Acad Sci U S A 111: 8299-8304

Pubmed: [Author and Title](#)

Downloaded from www.plantphysiol.org on January 19, 2017 - Published by www.plantphysiol.org
Copyright © 2017 American Society of Plant Biologists. All rights reserved.

CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Reggiori F, Wang C, Stromhaug P, Shintani T, Klionsky D (2003) Vps51 is part of the yeast Vps fifty-three tethering complex essential for retrograde traffic from the early endosome and Cvt vesicle completion. J Biol Chem 278: 5009-5020

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Rodriguez PA, Bos JI (2013) Toward understanding the role of aphid effectors in plant infestation. Mol Plant Microbe Interact 26: 25-30

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Rodriguez PA, Stam R, Warbroek T, Bos JIB (2014) Mp10 and Mp42 from the aphid species Myzus persicae trigger plant defenses in Nicotiana benthamiana through different activities. MPMI 27: 30-39

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Sandström J, Telang A, Moran NA (2000) Nutritional enhancement of host plants by aphids - a comparison of three aphid species on grasses. J Insect Physiol. 46: 33-40

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Schindler C, Chen Y, Pu J, Guo X, Bonifacino JS (2015) EARP is a multisubunit tethering complex involved in endocytic recycling. Nat Cell Biol 17: 639-650

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Schulze-Lefert P, Panstruga R (2011) A molecular evolutionary concept connecting nonhost resistance, pathogen host range, and pathogen speciation. Trends Plant Sci 16: 117-125

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Song J, Win J, Tian M, Schornack S, Kaschani F, Ilyas M, van der Hoorn RA, Kamoun S (2009) Apoplastic effectors secreted by two unrelated eukaryotic plant pathogens target the tomato defense protease Rcr3. Proc Natl Acad Sci U S A 106: 1654-1659

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Teh OK, Hofius D (2014) Membrane trafficking and autophagy in pathogen-triggered cell death and immunity. J Exp Bot 65: 1297-1312

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Thorpe P, Cock PJ, Bos J (2016) Comparative transcriptomics and proteomics of three different aphid species identifies core and diverse effector sets. BMC Genomics 17: 172

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Ustun S, Bartetzko V, Bornke F (2013) The Xanthomonas campestris type III effector XopJ targets the host cell proteasome to suppress salicylic-acid mediated plant defence. PLoS Pathog 9: e1003427

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zheng X, McLellan H, Fraiture M, Liu X, Boevink PC, Gilroy EM, Chen Y, Kandel K, Sessa G, Birch PR, Brunner F (2014) Functionally redundant RXLR effectors from Phytophthora infestans act at different steps to suppress early flg22-triggered immunity. PLoS Pathog 10: e1004057

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)